

Fisher-Symmetric Informationally Complete Measurements for Pure States

Nan Li,^{1,2,*} Christopher Ferrie,^{1,3,†} Jonathan A. Gross,^{1,‡} Amir Kalev,^{1,§} and Carlton M. Caves^{1,4,||}

¹*Center for Quantum Information and Control, University of New Mexico, Albuquerque, New Mexico 87131-0001, USA*

²*Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China*

³*Centre for Engineered Quantum Systems, School of Physics, The University of Sydney, Sydney, New South Wales 2006, Australia*

⁴*Centre for Engineered Quantum Systems, School of Mathematics and Physics, University of Queensland, Brisbane, Queensland 4072, Australia*

(Received 24 July 2015; published 5 May 2016)

We introduce a new kind of quantum measurement that is defined to be symmetric in the sense of uniform Fisher information across a set of parameters that uniquely represent pure quantum states in the neighborhood of a fiducial pure state. The measurement is locally informationally complete—i.e., it uniquely determines these parameters, as opposed to distinguishing two arbitrary quantum states—and it is maximal in the sense of a multiparameter quantum Cramér-Rao bound. For a d -dimensional quantum system, requiring only local informational completeness allows us to reduce the number of outcomes of the measurement from a minimum close to but below $4d - 3$, for the usual notion of global pure-state informational completeness, to $2d - 1$.

DOI: 10.1103/PhysRevLett.116.180402

A fundamental consequence of quantum mechanics is the inability to determine the quantum state of a single physical system. A well-posed problem in quantum state tomography is this: given many copies of a quantum system, all assumed to be in the same state ρ , find a measurement to perform on each copy that is sufficient to specify ρ uniquely in the limit of an infinite number of copies, i.e., from the outcome probabilities.

In quantum theory, measurements are represented by positive-operator-valued measures (POVMs), whose elements, E^ξ , are positive operators satisfying the completeness condition, $\sum_\xi E^\xi = \mathbb{1}$. If we perform the measurement on a system in state ρ , the probability of outcome ξ is $p^\xi = \text{tr}(\rho E^\xi)$. If the statistics of the outcome probabilities are sufficient to determine all the parameters of the state uniquely, then the POVM is said to be a tomographically or informationally complete POVM (IC-POVM).

In this Letter, we consider measurements whose outcome probabilities, though not globally informationally complete, can determine a quantum state in a local neighborhood of a fixed, but arbitrary fiducial state; we dub such measurements locally informationally complete. We quantify the information content of a measurement using the multiparameter Fisher-information matrix [1]. We look for measurements that satisfy two requirements. First, the information obtained should be maximal relative to a fundamental bound on the classical Fisher-information matrix, established by Gill and Massar [2]; by measuring the amount of classical Fisher information relative to the corresponding quantum Fisher information, the Gill-Massar (GM) bound is the multiparameter expression of the quantum Cramér-Rao bound [3–6]. Saturating the GM bound requires that the POVM elements be rank one.

Second, we look for measurements such that the classical Fisher information is distributed as uniformly as possible among the parameters of the quantum state. Measurements that satisfy these requirements are as efficient as possible for measuring all the parameters of a quantum state simultaneously; we call such measurements Fisher symmetric.

In this Letter, we specialize to pure states [7], where Fisher symmetry means that all the parameters of the pure state are determined with the same resolution relative to the corresponding quantum Cramér-Rao bound, and we refer to the measurements that meet our requirements as pure Fisher-symmetric informationally complete (PFSIC). We show that $2d - 1$ outcomes are necessary and, by example, sufficient for a PFSIC measurement, in contrast to a minimum close to, but below, $4d - 3$ outcomes required for global pure-state informational completeness [8]. Moreover, in accordance with the GM bound, each of the $2d - 2$ parameters of the pure state is determined with a resolution half that of a separate quantum-limited measurement of that parameter.

What is the minimal number of elements for a POVM to be globally informationally complete? A full-rank quantum state, described by a normalized density operator ρ in a d -dimensional Hilbert space, is specified by $d^2 - 1$ real parameters. Since the outcome probabilities establish a series of linear constraints for the mixed-state parameters and the POVM operators have to satisfy the completeness condition, it is simple to conclude that an IC-POVM must have at least d^2 elements. If, in addition, one asks for a minimal, rank-one IC-POVM and demands global symmetry in the geometry of the rank-one POVM elements, one arrives at SIC-POVMs [9–11], whose existence in all

dimensions is a topic of both mathematical and physical interest.

These considerations change if one knows that ρ is a pure state. Since the relation between the outcome probabilities and the pure-state parameters is quadratic rather than linear, the problem of the minimal number of elements in a global IC-POVM for pure states is more complicated. Flammia, Silberfarb, and Caves [12] considered pure-state informationally (PSI-)complete POVMs, whose outcome probabilities are sufficient to determine a generic pure state (up to a global phase), i.e., all states except for a set of pure states that is dense only on a set of measure zero. They showed that the minimal number of elements for these POVMs is $2d$, and they conjectured that $2d$ outcomes suffice even if the POVM elements are rank one. Finkelstein [13] confirmed this conjecture, but went on to show that if a rank-one POVM achieves global informational completeness for all pure states, not just a generic set [he called such a POVM PSI really (PSIR-)complete], then the POVM must have at least $3d - 2$ POVM elements. Finkelstein left open the question of whether a PSIR-complete POVM with this number of outcomes exists. Recently, in a *tour de force* of mathematical physics, Heinosaari, Mazzarella, and Wolf [8] showed that a POVM that identifies all pure states has minimally $4d - 3 - c(d)\alpha(d)$ outcomes, where $1 \leq c(d) \leq 2$ and $\alpha(d)$ is the number of 1's in the binary expansion of $d - 1$. This brief discussion illustrates the complicated nature of global informational completeness for pure states.

Here, we consider a different tomographic problem, which might be called local or point tomography. In this problem, an experimenter has more than just the promise of a pure state: she knows which pure state she is trying to prepare and knows she can prepare it quite well, except that the unitary operators used in the preparation have systematic errors; though the prepared state is pure, it is different from the fiducial state the experimenter is aiming for. The experimenter wants to find a measurement sufficient to identify all small discrepancies from this fiducial state. We quantify the “goodness” of a measurement with the multi-parameter Fisher-information matrix [1]. Fisher information is a key tool in statistics, which allows one to bound the performance of any estimator, and it has played a prominent role in the development of quantum information and metrology, where it has been suitably generalized to the quantum setting [3–6]. It is the appropriate tool here because it provides a definitive answer to questions about estimating local deviations from a fiducial state.

Now, we formulate the problem with precision, first generally for mixed states and, then, for the pure-state context analyzed in this Letter [7]. An unknown quantum state $\rho(\mathbf{x})$ depends on a vector $\mathbf{x} = (x^1, \dots, x^p)$ of p real parameters. For density operators of rank ℓ in $d \geq \ell$ dimensions, $p = 2d\ell - \ell^2 - 1$; for full-rank density operators, $p = d^2 - 1$, and for pure states, $p = 2d - 2$. The fiducial state can be labeled by $\mathbf{x} = \mathbf{0}$. In the following,

we are interested in the quantum and classical Fisher-information matrices evaluated at the fiducial state, i.e., at $\mathbf{x} = \mathbf{0}$.

The quantum Fisher information is a $p \times p$ real, symmetric matrix $\mathbf{Q}(\rho)$, whose matrix elements are

$$Q_{\alpha\beta}(\rho) = \frac{1}{2} \text{tr}[\rho(L_\alpha L_\beta + L_\beta L_\alpha)], \quad (1)$$

with the (Hermitian) symmetric logarithmic derivative (SLD) operators L_α , one for each parameter, determined implicitly by $\partial\rho/\partial x^\alpha = \frac{1}{2}(L_\alpha\rho + \rho L_\alpha)$. We can reparametrize the quantum state to make \mathbf{Q} the identity matrix.

Given a POVM, with elements E^ξ and outcome probabilities $p^\xi = \text{tr}(\rho E^\xi)$, the classical Fisher-information matrix is a $p \times p$ real, symmetric matrix \mathbf{C} , defined by

$$C_{\alpha\beta} = \sum_{\xi} \frac{1}{p^\xi} \frac{\partial p^\xi}{\partial x^\alpha} \frac{\partial p^\xi}{\partial x^\beta}. \quad (2)$$

The sum over ξ here is restricted to E^ξ that are not orthogonal to the fiducial state ρ , i.e., for which $p^\xi = \text{tr}(\rho E^\xi) \neq 0$.

Gill and Massar [2] proved that for any states $\rho(\mathbf{x})$ and any POVM, the classical Fisher-information matrix \mathbf{C} satisfies

$$\text{tr}(\mathbf{Q}^{-1}\mathbf{C}) = \text{tr}(\mathbf{Q}^{-1/2}\mathbf{C}\mathbf{Q}^{-1/2}) \leq d - 1, \quad (3)$$

with equality if and only if all the POVM elements are rank one, and none is orthogonal to the fiducial state. The GM quantity, $\text{tr}(\mathbf{Q}^{-1}\mathbf{C})$, is invariant under reparametrization of the quantum states. It is most easily interpreted when the parameters are chosen so that \mathbf{Q} is the identity matrix. The unit elements on the diagonal of \mathbf{Q} then express the quantum limit, called the quantum Cramér-Rao bound, on estimating each of the parameters separately [6]. The corresponding diagonal elements of the classical Fisher matrix give the performance of the POVM in determining these same parameters in units of the quantum limit. The GM quantity is the sum of the diagonal elements of \mathbf{C} ; the bound (3) expresses the quantum limit on estimating all the parameters simultaneously and, as such, is a Fisher-information expression of the uncertainty principle. Zhu [14] has made use of the positive symmetric matrix $\mathbf{Q}^{-1/2}\mathbf{C}\mathbf{Q}^{-1/2}$ in a study of information complementarity and incompatible observables.

We call a POVM Fisher symmetric if it saturates the GM bound (3) and has a classical Fisher matrix distributed as uniformly as possible among the parameters of the quantum state. What we mean by as uniformly as possible is that the measurement minimizes the quadratic quantity

$$\begin{aligned}\text{tr}(\mathbf{Q}^{-1}\mathbf{C}\mathbf{Q}^{-1}\mathbf{C}) &= \text{tr}((\mathbf{Q}^{-1/2}\mathbf{C}\mathbf{Q}^{-1/2})^2) \\ &\geq \frac{[\text{tr}(\mathbf{Q}^{-1/2}\mathbf{C}\mathbf{Q}^{-1/2})]^2}{p} \\ &= \frac{(d-1)^2}{p}.\end{aligned}\quad (4)$$

The inequality follows directly from minimizing this quadratic quantity subject to the linear trace constraint. The absolute minimum is achieved if and only if the classical Fisher matrix is proportional to the quantum Fisher matrix, i.e., $\mathbf{C} = (d-1)\mathbf{Q}/p$. For full-rank density operators, however, there is generally no POVM that achieves the absolute minimum; it can be achieved only for qubits and for the maximally mixed state in all dimensions [7]. For any density operator, however, the classical Fisher matrices are a convex set under coin-flipping convex combinations of the underlying POVMs; since the quantity (4) is convex, global minima are guaranteed to exist. Thus, the study of Fisher symmetry for full-rank density operators will be focused on finding what the minimum value is and what POVMs achieve it; such POVMs determine a quantum state locally as efficiently as possible.

For pure states, it is possible to achieve $\mathbf{C} = \frac{1}{2}\mathbf{Q}$, as we now show. A POVM that achieves $\mathbf{C} = \frac{1}{2}\mathbf{Q}$, what we call a PFSIC, can estimate all the parameters of a pure state with half the quantum-limited resolution with which each could be estimated separately. The unknown pure state $\rho(\mathbf{x}) = |\Psi(\mathbf{x})\rangle\langle\Psi(\mathbf{x})|$ depends on a vector \mathbf{x} of $2d-2$ real parameters. We denote the fiducial state $\mathbf{x} = \mathbf{0}$ by $|0\rangle$, i.e., $|\Psi(\mathbf{0})\rangle = |0\rangle$. By assumption, the unknown pure state $|\Psi(\mathbf{x})\rangle$ is close to $|0\rangle$ and, thus, can be parametrized to linear order as

$$|\Psi(\mathbf{x})\rangle = |0\rangle + \sum_{k=1}^{d-1} (x^{k0} + ix^{k1})|k\rangle, \quad (5)$$

where the set $\{|k\rangle\}_{k=0,1,\dots,d-1}$ is an orthonormal basis, $\mathbf{x} = (x^{1,0}, x^{1,1}, \dots, x^{d-1,0}, x^{d-1,1})$, and $|x^{k\sigma}| \ll 1$ for $k = 1, \dots, d-1$, $\sigma = 0, 1$. Keeping only the terms linear in the parameters, we get

$$\rho(\mathbf{x}) = |0\rangle\langle 0| + \sum_{k,\sigma} x^{k\sigma} X_{k\sigma} = |0\rangle\langle 0| + \sum_{\alpha} x^{\alpha} X_{\alpha}, \quad (6)$$

where

$$X_{k\sigma} = (-i)^{\sigma} (|0\rangle\langle k| + (-1)^{\sigma}|k\rangle\langle 0|), \quad (7)$$

i.e., $X_{k0} = |0\rangle\langle k| + |k\rangle\langle 0|$ and $X_{k1} = -i|0\rangle\langle k| + i|k\rangle\langle 0|$, for $k = 1, \dots, d-1$. In Eq. (6), we ignore second-order terms because they do not contribute either to the quantum Fisher information or to the measurement-induced classical Fisher information at the fiducial point. In accordance with

the foregoing, we sometimes let a single Greek index stand for both k and σ , as in the last form of Eq. (6). The Hermitian operators X_{α} satisfy $\text{tr}(X_{\alpha}X_{\beta}) = 2\delta_{\alpha\beta}$.

When $\rho(\mathbf{x})$ is a pure state, the SLDs are easy to find. At the fiducial state, the SLDs are $L_{\alpha} = 2\partial\rho(\mathbf{x})/\partial x^{\alpha}|_{\mathbf{x}=\mathbf{0}} = 2X_{\alpha}$, and the quantum Fisher-information matrix is $\mathbf{Q} = 4\mathbf{I}_{2d-2}$, where \mathbf{I}_n denotes the $n \times n$ identity matrix. Thus, we have chosen from the start a parametrization that is only a uniform rescaling away from making the quantum Fisher matrix the identity.

Now, consider any POVM that saturates the GM bound, i.e., has n rank-one POVM elements, none of which is orthogonal to $\rho(\mathbf{0}) = |0\rangle\langle 0|$, and take the POVM elements to be

$$E^{\xi} = |\psi^{\xi}\rangle\langle\psi^{\xi}| = \sum_{k,j=0}^{d-1} a_k^{\xi} (a_j^{\xi})^* |k\rangle\langle j|, \quad (8)$$

where the POVM vectors are

$$|\psi^{\xi}\rangle = \sum_{k=0}^{d-1} a_k^{\xi} |k\rangle, \quad \xi = 0, \dots, n-1. \quad (9)$$

The POVM completeness condition, $\sum_{\xi} E^{\xi} = \mathbb{1}$, becomes $\sum_{\xi} (a_j^{\xi})^* a_k^{\xi} = \delta_{jk}$. Defining

$$a_k^{\xi} = b_k^{\xi} + ic_k^{\xi}, \quad k = 0, \dots, d-1, \quad (10)$$

and gathering up the various components into n -dimensional column vectors,

$$\mathbf{b}_k = \begin{pmatrix} b_k^0 \\ b_k^1 \\ \vdots \\ b_k^{n-1} \end{pmatrix}, \quad \mathbf{c}_k = \begin{pmatrix} c_k^0 \\ c_k^1 \\ \vdots \\ c_k^{n-1} \end{pmatrix}, \quad k = 0, 1, \dots, d-1, \quad (11)$$

we can put the completeness condition in the form

$$\begin{aligned}\mathbf{b}_j \cdot \mathbf{b}_k + \mathbf{c}_j \cdot \mathbf{c}_k &= \delta_{jk}, \\ \mathbf{b}_j \cdot \mathbf{c}_k - \mathbf{c}_j \cdot \mathbf{b}_k &= 0,\end{aligned} \quad j, k = 0, 1, \dots, d-1. \quad (12)$$

Choice of phase of the POVM vectors (9) allows us to make $a_0^{\xi} = b_0^{\xi}$ real and nonnegative (thus, $c_0^{\xi} = 0$) for all ξ . Since we now have $\mathbf{c}_0 = 0$, there are $2d-1$ nonzero vectors (11). It is useful to spell out, separately, the $j=0$ or $k=0$ parts of the completeness conditions,

$$\begin{aligned} \mathbf{b}_0 \cdot \mathbf{b}_0 &= 1, \\ \mathbf{b}_0 \cdot \mathbf{b}_k &= \mathbf{b}_0 \cdot \mathbf{c}_k = 0, \quad k = 1, \dots, d-1. \end{aligned} \quad (13)$$

The probability to obtain the outcome ξ at the fiducial state is

$$p^\xi = \langle 0|E^\xi|0\rangle = |a_0^\xi|^2 = (b_0^\xi)^2 > 0. \quad (14)$$

Notice that the vector \mathbf{b}_0 is a normalized vector that has strictly positive components. We also have $\partial p^\xi(\mathbf{x})/\partial x^\alpha = \text{tr}(E^\xi X_\alpha)$, i.e.,

$$\frac{\partial p^\xi}{\partial x^{k0}} = \langle k|E^\xi|0\rangle + \langle 0|E^\xi|k\rangle = 2b_0^\xi b_k^\xi, \quad (15a)$$

$$\frac{\partial p^\xi}{\partial x^{k1}} = -i\langle k|E^\xi|0\rangle + i\langle 0|E^\xi|k\rangle = 2b_0^\xi c_k^\xi. \quad (15b)$$

The definition (2) of the classical Fisher matrix gives, for $j, k = 1, \dots, d-1$,

$$C_{j0,k0} = 4 \sum_{\xi} b_j^\xi b_k^\xi = 4\mathbf{b}_j \cdot \mathbf{b}_k, \quad (16a)$$

$$C_{j1,k1} = 4 \sum_{\xi} c_j^\xi c_k^\xi = 4\mathbf{c}_j \cdot \mathbf{c}_k, \quad (16b)$$

$$C_{j0,k1} = C_{k1,j0} = 4 \sum_{\xi} b_j^\xi c_k^\xi = 4\mathbf{b}_j \cdot \mathbf{c}_k. \quad (16c)$$

The classical Fisher matrix is a matrix of inner products of the $2d-2$ n -dimensional vectors $\{2\mathbf{b}_k, 2\mathbf{c}_k\}_{k=1}^{d-1}$. The rank of such a matrix, called a Gram matrix, is the dimension of the span of the vectors going into the inner products, so the rank of the classical Fisher matrix is bounded above by $\min(n, 2d-2)$.

In the pure-state case, the Fisher-symmetry condition becomes $\mathbf{C} = \frac{1}{2}\mathbf{Q} = 2\mathbf{I}_{2d-2}$. To satisfy this condition, \mathbf{C} must be full rank, i.e., have rank $2d-2$, which implies that $n \geq 2d-2$. Since \mathbf{C} is full rank, it is invertible, and the measurement is locally informationally complete in that it uniquely determines all parameters in the limit of infinitely many measurements.

In the parametrization we are using, the Fisher-symmetric POVM must satisfy

$$\begin{aligned} \frac{1}{2}C_{j0,k0} &= 2\mathbf{b}_j \cdot \mathbf{b}_k = \delta_{jk}, \\ \frac{1}{2}C_{j1,k1} &= 2\mathbf{c}_j \cdot \mathbf{c}_k = \delta_{jk}, \quad j, k = 1, \dots, d-1. \\ \frac{1}{2}C_{j0,k1} &= 2\mathbf{b}_j \cdot \mathbf{c}_k = 0, \end{aligned} \quad (17)$$

Combining these Fisher-symmetry conditions with the completeness conditions (13), we see that the vectors $\{\sqrt{2}\mathbf{b}_k, \sqrt{2}\mathbf{c}_k\}_{k=1}^{d-1}$ are a set of $2d-2$ orthonormal vectors

in an n -dimensional subspace, which is orthogonal to \mathbf{b}_0 . We can now conclude that $n \geq 2d-1$.

It is clear that PFSICs exist for all $n \geq 2d-1$, since they can be constructed by choosing an n -dimensional vector \mathbf{b}_0 with all positive components and then finding $2d-2$ orthonormal vectors in the subspace of dimension $n-1 \geq 2d-2$ orthogonal to \mathbf{b}_0 . In the Supplemental Material [15], we construct a minimal ($n = 2d-1$) PFSIC POVM by choosing \mathbf{b}_0 symmetrically, i.e., $\mathbf{b}_0^T = (1, 1, \dots, 1)/\sqrt{2d-1}$ and using a manifestly symmetric method to construct the remaining vectors. For qubits, this PFSIC reduces to the trine measurement, with the POVM vectors corresponding to outcomes in the equatorial plane of the Bloch sphere. The trine measurement is locally informationally complete for pure states near the north pole of the Bloch sphere, but it is not globally informationally complete because it cannot distinguish states whose Bloch vectors differ by a sign flip of the z component of the Bloch vector. This example illustrates how PFSICs fail to be globally informationally complete. Yet, they are informationally complete over a region containing the fiducial state: here, the northern hemisphere of the Bloch sphere, a more complicated shape in higher dimensions.

In the Supplemental Material [15], we also construct a POVM that comes from flipping a coin with probabilities p_χ and p_τ to choose between measuring in one of two orthonormal bases

$$\begin{aligned} |\chi^\xi\rangle &= u_0^\xi|0\rangle + \sum_{j=1}^{d-1} u_j^\xi|j\rangle, \\ |\tau^\xi\rangle &= -iu_0^\xi|0\rangle + \sum_{j=1}^{d-1} u_j^\xi|j\rangle, \end{aligned} \quad \xi = 0, 1, \dots, d-1, \quad (18)$$

with $u_0^\xi > 0$. With the POVM vectors chosen to be $|\psi^\xi\rangle = \sqrt{p_\chi}|\chi^\xi\rangle$ and $|\psi^{d+\xi}\rangle = \sqrt{p_\tau}|\tau^\xi\rangle$, the Fisher-information matrix (16) is diagonal, with diagonal components $C_{j0,j0} = 4p_\chi$ for the estimates of the real parts, x^{k0} , of the amplitudes, and $C_{j1,k1} = 4p_\tau$ for estimates of the imaginary parts, x^{k1} . When $p_\chi = p_\tau = \frac{1}{2}$, we have a PFSIC. Other weightings of the coin give different trade-offs, within the GM bound, between determining the real and imaginary parts of the amplitudes. This example illustrates the sense in which the GM bound can be thought of as an uncertainty principle for measuring, simultaneously, the parameters that specify a pure state. For a qubit, the two bases (18) correspond to measurements of the Pauli operators σ_x and σ_y .

It is useful to clarify what freedom we have in choosing a PFSIC. Starting with a minimal PFSIC, i.e., with an orthonormal set $\{\mathbf{b}_0, \sqrt{2}\mathbf{b}_k, \sqrt{2}\mathbf{c}_k\}_{k=1, \dots, d-1}$, of $(2d-1)$ -dimensional real vectors, where \mathbf{b}_0 has all positive components, we can do any (active) orthogonal transformation O to get a new set $\mathbf{B}_0 = O\mathbf{b}_0$ and $\mathbf{B}_j = O\mathbf{b}_j$, $\mathbf{C}_j = O\mathbf{c}_j$, for $j = 1, \dots, d-1$, subject to the requirement that $B_0^\xi > 0$ for all ξ [16]. To get a nonminimal PFSIC, we add

additional dimensions to the real vector space and allow O to map into these extra dimensions. In terms of components, we have

$$B_j^\xi = \sum_{\eta=0}^{2d-2} O_\eta^\xi b_j^\eta, \quad j = 0, 1, \dots, d-1, \quad (19)$$

$$C_j^\xi = \sum_{\eta=1}^{2d-2} O_\eta^\xi c_j^\eta, \quad j = 1, \dots, d-1. \quad (20)$$

Letting $|\phi^\xi\rangle$ be the POVM vectors for the primed real vectors, we have

$$|\phi^\xi\rangle = \sum_{j=0}^{d-1} (B_j^\xi + iC_j^\xi)|j\rangle = \sum_{\eta=0}^{2d-2} O_\eta^\xi |\psi^\eta\rangle. \quad (21)$$

Thus, our freedom is to do any orthogonal mixing of the POVM vectors, subject to $B_0^\xi = \langle 0|\phi^\xi\rangle > 0$ [16].

How does this compare with the usual freedom for rank-one POVMs? In complete analogy with the Hughston-Josza-Wootters freedom for pure-state ensemble decompositions of a density operator [17], for POVMs, we are asking for the freedom in writing the unit operator as a sum of rank-one operators. Generally, that freedom is the ability to mix the POVM vectors with any unitary matrix, which always yields another POVM. The restriction here is that we can only use real unitaries, i.e., orthogonal matrices, that leave \mathbf{b}_0 with all nonzero components.

The most complete information we can have of a physical system is its quantum state. We access this information by making repeated measurements on systems prepared in the same state. There are two fundamental and practical questions about such measurements: (i) Which schemes are sufficient to specify the state uniquely? (ii) Which schemes provide the most information per measurement? Typically, such questions are studied separately, the former using tools of linear algebra and convex geometry and the latter using quantum generalizations of Fisher statistics or other methods. Here, we bring these two lines of questioning together to construct measurements that are minimal, symmetric, and informationally complete, but in a local, Fisher-statistical sense rather than in the global, geometric sense. We have explicitly constructed measurements with $2d-1$ outcomes that are sufficient to estimate, simultaneously, all the parameters of pure quantum states near a fiducial state. Moreover, these measurements provide equal and optimal information about all the parameters. The quantum price for the simultaneous estimation, in accordance with the GM uncertainty principle (3), is that each parameter is determined with half the sensitivity with which it could be determined separately.

This work was supported in part by US National Science Foundation Grants No. PHY-1212445 and

No. PHY-1521016 and by US Office of Naval Research Grant No. N00014-15-1-2167. N. L. was also supported by the National Natural Science Foundation of China (Grants No. 11405262, No. 61134008, and No. 11375259), the National Center for Mathematics and Interdisciplinary Sciences, Chinese Academy of Sciences (Grant No. Y029152K51), and the Key Laboratory of Random Complex Structures and Data Science, Chinese Academy of Sciences (Grant No. 2008DP173182). C. F. was also supported by the Canadian Government through the NSERC PDF program, the IARPA MQCO program, the Australian Research Council via EQUs Project No. CE11001013, and U.S. Army Research Office Grants No. W911NF-14-1-0098 and No. W911NF-14-1-0103.

*linan@amss.ac.cn

†csferrie@gmail.com

‡jagross@unm.edu

§amirk@unm.edu

||ccaves@unm.edu

- [1] R. A. Fisher, *Proc. Cambridge Philos. Soc.* **22**, 700 (1925).
- [2] R. D. Gill and S. Massar, *Phys. Rev. A* **61**, 042312 (2000).
- [3] H. P. Yuen and M. Lax, *IEEE Trans. Inf. Theory* **19**, 740 (1973).
- [4] C. W. Helstrom, *Quantum Detection and Estimation Theory* (Academic, New York, 1976).
- [5] A. S. Holevo, *Probabilistic and Statistical Aspects of Quantum Theory* (North-Holland, Amsterdam, 1982).
- [6] S. L. Braunstein and C. M. Caves, *Phys. Rev. Lett.* **72**, 3439 (1994).
- [7] In subsequent papers, we develop the theory of Fisher-symmetric informationally complete measurements for mixed states.
- [8] T. Heinosaari, L. Mazzarella, and M. M. Wolf, *Commun. Math. Phys.* **318**, 355 (2013).
- [9] G. Zauner, Ph.D. thesis, University of Vienna, 1999, <https://www.mat.univie.ac.at/~neum/ms/zauner.pdf> [*Int. J. Quantum. Inform.* **09**, 445 (2011)].
- [10] J. M. Renes, R. Blume-Kohout, A. J. Scott, and C. M. Caves, *J. Math. Phys.* **45**, 2171 (2004).
- [11] D. M. Appleby, H. B. Dang, and Christopher A. Fuchs, *Entropy* **16**, 1484 (2014).
- [12] S. T. Flammia, A. Silberfarb, and C. M. Caves, *Found. Phys.* **35**, 1985 (2005).
- [13] J. Finkelstein, *Phys. Rev. A* **70**, 052107 (2004).
- [14] H. Zhu, *Sci. Rep.* **5**, 14317 (2015).
- [15] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.116.180402> for an explicit construction of two kinds of PFSICs: minimal ($2d-1$ outcomes); minimal-plus-one ($2d$ outcomes), which consists of coin flipping between measurements in two orthogonal bases.
- [16] We can start with the requirement that $B_0^\xi = \langle 0|\phi^\xi\rangle \neq 0$ and make these components positive by reflections in the real vector space (equivalently, by changing the sign of the relevant POVM vectors $|\phi^\xi\rangle$).
- [17] L. P. Hughston, R. Josza, and W. K. Wootters, *Phys. Lett. A* **183**, 14 (1993).